

The Hip's Influence on Low Back Pain: A Distal Link to a Proximal Problem

Michael P. Reiman, P. Cody Weisbach, and Paul E. Glynn

Low back pain (LBP) is a multifactorial dysfunction, with one of the potential contributing factors being the hip joint. Currently, research investigating the examination and conservative treatment of LBP has focused primarily on the lumbar spine. The objective of this clinical commentary is to discuss the potential link between hip impairments and LBP using current best evidence and the concept of regional interdependence as tools to guide decision making and offer ideas for future research.

Keywords: strength, rehabilitation

In day-to-day clinical practice it is often difficult to identify the source of symptoms in patients with low back pain (LBP).¹ Abenhaim et al² noted that a small percentage of individuals with LBP have an identifiable pathoanatomical source. Further clouding the picture are multiple studies indicating the potential inability of diagnostic imaging to identify the pain source, influence prognosis, or affect outcomes.³⁻⁶ Research has demonstrated the effectiveness of subgrouping patients into a classification system based on signs and symptoms indicating their likelihood to respond to specific treatments. This classification approach has produced improved outcomes and high levels of reliability as compared with clinical-practice guidelines.⁷⁻⁹ For treating clinicians, these findings help guide decision making and improve results; however, not all patients will fit into a treatment-based subgroup. The treating therapist must then rely on an impairment-based approach, identifying potential local or remote contributors to the patient's area of primary concern. Recent advances in research have begun to indicate the importance of this regional approach to musculoskeletal examination, but the contributing areas remote to the lumbar spine have yet to be identified.

Given the biomechanical relationship between the hip and the low back, specifically the **multiple shared muscles (psoas, quadratus lumborum, erector spinae, gluteus maximus, etc)**, one must consider the hip joint a potential contributor to LBP. Contraction of these muscles will affect motion at the spine, pelvis, and hip because of common attachment sites. It is generally understood that movement at one of these areas will necessitate compensatory movement at the others because

Reiman is with the Physical Therapy Dept, Wichita State University, Wichita, KS 67260-0043. Weisbach is a staff physical therapist, and Glynn, the supervisor of staff development, at Newton-Wellesley Hospital, Newton, MA.

of these common muscle attachments and the concept of regional interdependence.¹⁰ By understanding the association between the hip and low back, clinicians might gain insight into the management of this challenging patient population.

Recently, clinical prediction rules (CPRs) have begun to help clinicians and have been gaining increased attention in the rehabilitation literature. CPRs are tools that help identify characteristics of individuals who might benefit from specific interventions. Two such rules have been developed pertaining to individuals with LBP that have helped shape the treatment-based classification system. Flynn et al¹¹ developed a CPR to identify individuals with acute LBP likely to demonstrate short-term benefit from lumbopelvic manipulation, which was later validated.¹² Five factors associated with success were identified for this intervention.^{11,12} Similarly, Hicks et al¹³ developed a CPR identifying individuals with LBP who might benefit from a lumbar-stabilization program. A common finding in both the manipulation and stabilization CPRs was the presence of adequate hip range of motion (ROM). In the manipulation CPR, patients with internal rotation greater than 35° in 1 or both hips had an increased likelihood of benefiting from manipulation of the lumbopelvic region.¹¹ In the lumbar-stabilization CPR, patients with a straight-leg raise greater than 91° were more likely to experience reduced disability at the 8-week follow-up.¹³ Prediction rules, as outlined here, are meant to classify patients to specific treatments; their purpose is not to explain how or why the variables influence the outcome. Therefore, a full understanding of the relationship between the hip and lumbar spine has yet to be elucidated. The concept of regional interdependence might help explain why hip characteristics influence the classification of individuals with LBP.

Regional interdependence as it applies to musculoskeletal physical therapy refers to “the concept that seemingly unrelated impairments in a remote anatomical region may contribute to, or be associated with the patient’s primary complaint.”^{10(p658)} Examples in the rehabilitative literature include manipulation of the thoracic spine for mechanical neck pain,¹⁴ lateral epicondylalgia,¹⁵ or shoulder dysfunction.^{16,17} Likewise, joint mobilization and strengthening treatments of the hip joint have been advocated for knee osteoarthritis,^{18–20} and joint manipulation for the low back has been advocated for knee^{21–23} and hip impairments.²⁴ Currently, manual therapy treatment of the hip joint, including mobilization, manipulation, and stretching, for LBP has only been investigated in a case report²⁵ and in the spinal-stenosis population^{26,27}; however, existing evidence does suggest a relationship between LBP and decreased hip mobility^{28–34} and/or strength.^{35–39}

Hip–Spine Syndrome

The concept of a biomechanical link between the hip joint and the lumbar spine has been described as hip–spine syndrome (HSS).⁴⁰ HSS specifically depicts the influence of a pathological hip joint on the alignment of the spine and subsequent muscle length and joint forces.⁴⁰

The most recent documentation of this relationship has been that of severe hip osteoarthritis (OA) potentially causing abnormal spinal sagittal alignment and ensuing LBP.⁴¹ Ben-Galim et al⁴¹ evaluated the effects of surgical treatment of hip

OA on low back disability in patients preoperative and postoperative total hip replacement and found significant ($P < .01$) improvements in both visual analog scores for LBP and Oswestry Disability Index scores after surgery that remained at the 2-year follow-up.

Other specific related interactions of the hip and spine in HSS can include a hip-flexion contracture resulting in compensatory hyperlordosis of the lumbar spine or a posteriorly inclined pelvis with increased kyphotic posture and primary or rapidly destructive hip OA.^{40,42–48} In each of these examples, although there is a relationship between the hip and spine, the evidence demonstrating the significance of its effect on LBP is deficient.

Although the biomechanical influences of the hip on LBP are not fully evident at this time, the current level of evidence does support a regional relationship between the 2 areas. From the preliminary work of Ben-Galim et al,⁴¹ one can begin to appreciate the importance of further investigating hip ROM, as well as regional soft-tissue characteristics, in patients with LBP.

Hip ROM and LBP

The proposed regional relationship between hip ROM and LBP based on the preliminary work noted in HSS has been further substantiated through numerous studies. Ellison et al²⁸ compared the hip rotation of patients with LBP with that of unmatched controls without LBP. Patients with LBP more frequently demonstrated asymmetrical hip-rotation ROM, with internal rotation (IR) being less than external rotation (ER). Chesworth et al²⁹ also compared hip-rotation ROM in subjects with LBP with a control group matched for age, gender, height, and weight. Both IR and ER were significantly limited in the LBP group compared with the control group. Cibulka et al³⁰ observed subjects with LBP to have bilateral asymmetries in hip IR, whereas patients with LBP and sacroiliac-joint dysfunction had unilateral asymmetries. A cross-sectional study by Vad et al³² demonstrated that symptomatic, as compared with asymptomatic, professional golfers had significantly decreased hip ROM in a combination of flexion, abduction, and external rotation (FABER) of the lower extremity that leads in the golf swing when compared with the opposite leg. In addition, these same symptomatic golfers had significantly decreased hip IR ROM of the lead leg compared with the nonlead leg (11.8° vs 16.9°, respectively). No hip-ROM asymmetries were reported in the asymptomatic subjects. The aforementioned studies support the idea that ROM restrictions, whether pathological in the case of hip OA or nonpathological in the case of soft-tissue restrictions, are correlated with LBP.

Hip-Muscle Performance and LBP

The work investigating the effects of hip strength on LBP has focused primarily on athletes. In 2000 Nadler et al³⁵ examined the relationship of hip-extension and -abduction strength in athletes and their effect on future LBP and lower extremity injury. Female athletes with a history of LBP demonstrated a significant side-to-side asymmetry in hip-extensor strength.³⁵ No difference was found in hip-exten-

tor strength for male athletes with or without a history of injury or in hip-abduction-strength differences for either gender. In 2001 Nadler et al³⁶ examined hip-abductor and -extensor strength in college athletes. Logistic-regression analysis indicated a difference in side-to-side hip-extension strength as a potential predictive variable of future treatment for LBP among female athletes only.

Kankaanpää et al³⁸ investigated hip- and back-extension fatigability in subjects with chronic LBP. Using EMG spectral analysis, they reported that paraspinal fatigability was similar between groups, whereas the *gluteus maximus* fatigued more rapidly in the chronic LBP group. In addition, in a multifactorial cross-sectional study of 600 subjects with LBP, it was determined that hip-flexor, hip-abductor, and abdominal-muscle fatigability had a significant association with LBP.³⁹ The cited research indicates the evolving body of evidence linking strength decrements of the hip region to LBP.

Table 1 Level of Evidence Supporting Relationships Between Specific Hip and Low Back Dysfunction

Hip and low back relationship	Studies	Level of evidence supporting relationship
Hip–spine syndrome	Officerski & McNab ⁴⁰	Level III
	Ben-Galim et al ⁴¹	Level II
	Murata et al ⁴²	Level III
	Nakamura et al ⁴³	Level III
	Yoshimoto et al ⁴⁴	Level III
	Takemitsu et al ⁴⁵	Level III
	Sato et al ⁴⁶	Level III
	Itoi ⁴⁷	Level II
Watanabe et al ⁴⁸	Level III	
Hip range of motion and low back pain	Ellison et al ²⁸	Level III
	Chesworth et al ²⁹	Level III
	Cibulka et al ³⁰	Level III
	Sjolie ³¹	Level III
	Vad et al ³²	Level III
	Coplan ³³	Level III
	Mellin ³⁴	Level III
Hip-muscle performance and low back pain	Nadler et al ³⁵	Level III
	Nadler et al ³⁶	Level III
	Nadler et al ³⁷	Level III
	Kandaanpää et al ³⁸	Level III
	Nourbakhsh & Arab ³⁹	Level III

Recommendations

Although there is a lack of strong evidence (Table 1) with respect to the efficacy of treating the hip for LBP, this relationship still warrants consideration. Given the association of impaired hip ROM and LBP, treatment of hip-ROM deficits might be a beneficial treatment strategy. There have been studies that support the ability to improve hip-ROM deficits in patients with hip^{19,49} and knee²⁰ OA. MacDonald et al⁴⁹ explored the effects of manual therapy and exercise in patients with hip OA (as classified by the American College of Rheumatology criteria⁵⁰) in a case-series format. Mobilizations included both thrust and nonthrust techniques directed at the hypomobile areas of the hip capsule as determined by the treating clinician. All 7 subjects experienced decreased hip-region pain, increased function, increased hip ROM (especially into flexion and IR), and a median improvement of total hip ROM (summation of flexion, extension, abduction, IR, and ER) of 82°. Hoeksma et al¹⁹ conducted a randomized controlled trial to investigate the difference between a stretching and long-axis thrust hip-mobilization group versus an exercise group in patients with hip OA. Results indicated that the hip-mobilization and stretching group achieved greater increases in hip flexion-extension and IR-ER ROM, as well as decreased pain and improved function and walking distance. Cliborne et al²⁰ investigated the short-term responses to hip mobilization of hip ROM and hip pain during provocative testing in individuals with knee OA. Manual therapy techniques were directed at either the anterior or posterior capsule of the hip, depending on where patients felt pain during provocative testing. Subjects demonstrated a significant ($P < .05$) increase in mean composite hip ROM (sum of hip flexion, functional squat, and FABER ROM) of 12.1° immediately after mobilization.

Evidence to support treating the hip for LBP is limited to a case study,²⁵ a case series,²⁷ and 1 randomized controlled trial.²⁶ Cibulka²⁵ described the case of a 35-year-old male with unilateral LBP diagnosed as sacroiliac dysfunction. The subject was found to have hip-ER asymmetry that was treated with an impairment-based stretching and strengthening program aimed at the hip, as well as the low back. Results indicated a 38% reduction in disability as measured by the Oswestry Disability Index, which was maintained at 1-year follow-up. Whitman et al²⁷ examined the effect of manual therapy and exercise applied to the lumbar spine, hip, and lower extremity in patients with lumbar spinal stenosis in a case series, as well as a randomized controlled trial.²⁶ In both studies, impairment-based manual therapy treatments of the hips and lumbar spine were applied with accompanying home exercises. Outcomes indicated positive functional improvements at both the short- and long-term follow-ups.

Given the role that hip-extensor strength and endurance,^{38,39} along with the role hip-abductor and -adductor muscles, plays in lateral stability of the pelvis,⁵¹ it is suggested that clinicians carefully examine the strength of these groups. Based on exam findings, clinicians can implement strengthening or stretching exercises with an emphasis on hip-extensor endurance given the findings of Kankaanpää,³⁸ as well as extensor strength given the work of Nadler et al.³⁵⁻³⁷ The effect of hip-musculature strengthening on LBP is an area that requires further research to determine whether increased hip strength leads to decreases in pain

and improvements in function. Research is also needed to determine more specifically whether there is a certain subset of patients with LBP that would benefit the most from such strengthening.

In addition to producing improvements in ROM, hip mobilizations have been found to improve hip-extensor⁵² and -abductor⁵³ strength in normal individuals. Specifically, patients treated with grade IV mobilizations addressing the anterior hip capsule demonstrated immediate improvement in gluteus maximus⁵² strength, and those treated with grade IV mobilizations addressing the inferior capsule demonstrated immediate improvement of hip-abductor⁵³ strength, compared with control groups. Although these studies were conducted on unimpaired individuals, their results might be relevant to those with hip-joint dysfunction, because it has been demonstrated that the muscles of the pelvic girdle most commonly affected with hip pathologies are the gluteal muscles.⁵⁴ This assumption has yet to be proven in the current literature, and future research is suggested before implementing it in treatment.

Conclusion

Best current evidence supports the link between impairments at the hip and LBP. Research suggests that decreased hip ROM, hip-extensor strength, and hip-adductor or -flexor endurance might contribute to pain in the lumbar area. Because of this emerging relationship, we suggest that hip-joint ROM, muscle performance, anatomical alignment, and mobility be considered during examination of patients with LBP.

Identifying hip impairments would lead to an impairment-based approach to treatment because current evidence has not identified the subgroup of LBP patients who would specifically benefit from treatment aimed at the hip. Because of the lack of high-quality research to help guide decision making, clinicians are left to intervene with impairment-level treatments. Interventions should focus on restoring hip ROM through both thrust and nonthrust mobilizations aimed at the areas of restriction. Treatments might also include both strength and endurance training of the identified impaired hip musculature.

Future research should further test the theoretical basis of treating the hip for LBP. Case studies or case series investigating the effects of hip mobilization or specific hip strengthening in LBP and HSS populations would be helpful in describing how clinicians use these techniques to treat this population. CPRs to identify subgroups of LBP and HSS patients who will benefit from specific interventions aimed at the hip could help clinicians decide when it is most appropriate to use these techniques. Most important to substantiate this relationship would be randomized controlled trials to determine whether treating the hip adds benefit to treatment of lumbar-spine impairments, which would also be necessary to validate the CPR.

Until such research is done to better direct treatment, it is recommended that clinicians consider taking a regional approach to the examination and treatment of LBP. Attention should be paid to the hip joint and its surrounding soft tissue, and interventions should be applied based on the impairments identified.

References

1. Fogel GR, Esses SI. Hip spine syndrome: management of coexisting radiculopathy and arthritis of the lower extremity. *Spine J.* 2003;3:238–241.
2. Abenheim L, Rossignol M, Gobeille D, Bonvalot Y, Fines P, Scott S. The prognostic consequences in the making of the initial medical diagnosis or work-related back injuries. *Spine.* 1995;20:791–795.
3. Borenstein DG, O'Mara JW, Boden SD, et al. The value of magnetic resonance imaging of the lumbar spine to predict low-back pain in asymptomatic subjects. *J Bone Joint Surg Am.* 2001;83(9):1306–1311.
4. Modic MT, Obuchowski NA, Ross JS, et al. Acute low back pain and radiculopathy: MRI imaging findings and their prognostic roll and effect on outcome. *Radiology.* 2005;237:597–604.
5. Carragee E, Alamin T, Cheng I, Franklin T, van den Haak E, Hurwitz E. Are first-time episodes of serious LBP associated with new MRI findings? *Spine J.* 2006;6:624–635.
6. Kleinstuck F, Dvorak J, Mannion AF. Are “structural abnormalities” on magnetic resonance imaging a contraindication to the successful conservative treatment of chronic nonspecific low back pain? *Spine.* 2006;31:2250–2257.
7. Brennan GP, Fritz JM, Hunter SJ, et al. Identifying subgroups of patients with acute/subacute “nonspecific” low back pain: results of a randomized clinical trial. *Spine.* 2006;31:623–631.
8. Fritz JM, Delitto A, Erhard RE. Comparison of classification-base physical therapy with therapy based on clinical practice guidelines for patients with acute low back pain: a randomized clinical trial. *Spine.* 2003;28:1363–1371,1372.
9. Fritz JM, Brennan GP, Clifford SN, et al. An examination of the reliability of a classification algorithm for subgrouping patients with low back pain. *Spine.* 2006;31:77–82.
10. Wainner RS, Whitman JM, Cleland JA, Flynn TW. Regional interdependence: a musculoskeletal examination model whose time has come. *J Orthop Sports Phys Ther.* 2007;37(11):658–660.
11. Flynn T, Fritz J, Whitman J, et al. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with spinal manipulation. *Spine.* 2002;27(24):2835–2843.
12. Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med.* 2004;141(12):920–928.
13. Hicks GE, Fritz JM, Delitto A, McGill SM. Preliminary development of a clinical prediction rule for determining which patients with low back pain will respond to a stabilization exercise program. *Arch Phys Med Rehabil.* 2005;86(9):1753–1762.
14. Cleland JA, Childs JD, Fritz JM, Whitman JM, Eberhart SL. Development of a clinical prediction rule for guiding treatment of a subgroup of patients with neck pain: use of thoracic spine manipulation, exercise, and patient education. *Phys Ther.* 2007;87(1):9–23.
15. Cleland JA, Whitman JM, Fritz JM. Effectiveness of manual physical therapy to the cervical spine in the management of lateral epicondylalgia: a retrospective analysis. *J Orthop Sports Phys Ther.* 2004;34(11):713–722 discussion 722-4.
16. Bang MD, Deyle GD. Comparison of supervised exercise with and without manual physical therapy for patients with shoulder impingement syndrome. *J Orthop Sports Phys Ther.* 2000;30(3):126–137.
17. Bergman GJ, Winters JC, Groenier KH, et al. Manipulative therapy in addition to usual medical care for patients with shoulder dysfunction and pain: a randomized, controlled trial. *Ann Intern Med.* 2004;141(6):432–439.

18. Currier LL, Froehlich PJ, Carow SD, et al. Development of a clinical prediction rule to identify patients with knee pain and clinical evidence of knee osteoarthritis who demonstrate a favorable short-term response to hip mobilization. *Phys Ther.* 2007;87(9):1106–1119.
19. Hoeksma HL, Dekker J, Ronday HK, et al. Comparison of manual therapy and exercise therapy in osteoarthritis of the hip: a randomized clinical trial. *Arthritis Rheum.* 2004;51(5):722–729.
20. Cliborne AV, Wainner RS, Rhon DI, et al. Clinical hip tests and a functional squat test in patients with knee osteoarthritis: reliability, prevalence of positive test findings, and short-term response to hip mobilization. *J Orthop Sports Phys Ther.* 2004;34(11):676–685.
21. Suter E, McMorland G, Herzog W, Bray R. Decrease in quadriceps inhibition after sacroiliac joint manipulation in patients with anterior knee pain. *J Manipulative Physiol Ther.* 1999;22(3):149–153.
22. Suter E, McMorland G, Herzog W, Bray R. Conservative lower back treatment reduces inhibition in knee-extensor muscles: a randomized controlled trial. *J Manipulative Physiol Ther.* 2000;23(2):76–80.
23. Iverson CA, Sutlive TG, Crowell MS, et al. Lumbopelvic manipulation for the treatment of patients with patellofemoral pain syndrome: development of a clinical prediction rule. *J Orthop Sports Phys Ther.* 2008;38(6):297–312.
24. Cibulka MT, Delitto A. A comparison of two different methods to treat hip pain in runners. *J Orthop Sports Phys Ther.* 1993;17(4):172–176.
25. Cibulka MT. Low back pain and its relation to the hip and foot. *J Orthop Sports Phys Ther.* 1999;29(10):595–601.
26. Whitman JM, Flynn TW, Childs JD, et al. A comparison between two physical therapy treatment programs for patients with lumbar spinal stenosis: a randomized clinical trial. *Spine.* 2006;31(22):2541–2549.
27. Whitman JM, Flynn TW, Fritz JM. Nonsurgical management of patients with lumbar spinal stenosis: a literature review and a case series of three patients managed with physical therapy. *Phys Med Rehabil Clin N Am.* 2003;14:77–101.
28. Ellison JB, Rose SJ, Sahrman SA. Patterns of hip rotation range of motion: a comparison between healthy subjects and patients with low back pain. *Phys Ther.* 1990;70(9):537–541.
29. Chesworth BM, Padfield BJ, Helewa A, Stitt LW. A comparison of hip mobility in patients with low back pain and matched healthy subjects. *Physiother Can.* 1994;46(4):267–274.
30. Cibulka MT, Sinacore DR, Cromer GS, Delitto A. Unilateral hip rotation range of motion asymmetry in patients with sacroiliac joint regional pain. *Spine.* 1998;23(9):1009–1015.
31. Sjolie AN. Low-back pain in adolescents is associated with poor hip mobility and high body mass index. *Scand J Med Sci Sports.* 2004;14(3):168–175.
32. Vad VB, Bhat AL, Basrai D, Gebeh A, Aspergren DD, Andrews JR. Low back pain in professional golfers: the role of associated hip and low back range-of-motion deficits. *Am J Sports Med.* 2004;32(2):494–497.
33. Coplan JA. Ballet dancer's turnout and its relationship to self-reported injury. *J Orthop Sports Phys Ther.* 2002;32(11):579–584.
34. Mellin G. Correlations of hip mobility with degree of back pain and lumbar spinal mobility in chronic low-back pain patients. *Spine.* 1988;13(6):668–670.
35. Nadler SF, Malanga GA, DePrince M, Stitik TP, Feinberg JH. The relationship between lower extremity injury, low back pain, and hip muscle strength in male and female collegiate athletes. *Clin J Sport Med.* 2000;10(2):89–97.
36. Nadler SF, Malanga GA, Feinberg JH, Prybicien M, Stitik TP, DePrince MS. Relationship between hip muscle imbalance and occurrence of low back pain in collegiate athletes: a prospective study. *Am J Phys Med Rehabil.* 2001;80(8):572–577.

37. Nadler SF, Malanga GA, Bartoli LA, Feinberg JH, Prybicien M, Deprince M. Hip muscle imbalance and low back pain in athletes: influence of core strengthening. *Med Sci Sports Exerc.* 2002;34(1):9–16.
38. Kankaanpää M, Taimela S, Laaksonen D, Hänninen O, Airaksinen O. Back and hip extensor fatigability in chronic low back pain patients and controls. *Arch Phys Med Rehabil.* 1998;79:412–417.
39. Nourbakhsh MR, Ara AM. Relationship between mechanical factors and incidence of low back pain. *J Orthop Sports Phys Ther.* 2002;32:447–460.
40. Offierski CM, McNab I. Hip-spine syndrome. *Spine.* 1983;8:316–321.
41. Ben-Galim P, Ben-Galim T, Rand N, et al. Hip-spine syndrome: the effect of total hip replacement surgery on low back pain in severe osteoarthritis of the hip. *Spine.* 2007;32(19):2099–2102.
42. Murata Y, Utsumi T, Hanaoka E, et al. Changes of the lumbar alignment in the same persons over a period of ten years. *Clin Orthop Surg.* 2002;37:1419–1422.
43. Nakamura N, Sugano N, Nishi T, et al. Scintigraphic image patterns in dysplastic coxarthrosis. *Acta Orthop Scand.* 2003;74:159–164.
44. Yoshimoto H, Shigenobu S, Masuda T, et al. Spinopelvic alignment in patients with osteoarthritis of the hip. *Spine.* 2005;30:1650–1657.
45. Takemitsu Y, Harada Y, Iwahara T, et al. Lumbar degenerative kyphosis. clinical, radiological and epidemiological studies. *Spine.* 1988;13:1317–1326.
46. Sato K, Itoi E, Kasama F. Abnormal posture associated with osteoporosis. *J Musculoskelet Syst.* 1989;2:1451–1462.
47. Itoi E. Roentgenographic analysis of posture in spinal osteoporosis. *Spine.* 1991;16:750–756.
48. Watanabe W, Sato K, Itoi E, et al. Posterior pelvic tilt in patients with decreased lumbar lordosis decreases acetabular femoral head covering. *Orthopedics.* 2002;25:321–324.
49. MacDonald CW, Whitman JM, Cleland JA, Smith M, Hoeksma HL. Clinical outcomes following manual physical therapy and exercise for hip osteoarthritis: a case series. *J Orthop Sports Phys Ther.* 2006;36(8):588–599.
50. Altman R, Alarcon G, Appelrouth D, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hip. *Arthritis Rheum.* 1991;34:505–514.
51. Lee D. *The Pelvic Girdle: An Approach to Examination and Treatment of the Lumbo-Pelvic-Hip Region.* 2nd ed. New York, NY: Churchill Livingstone; 2004.
52. Yerys S, Makofsky H, Byrd C, Pennachio J, Cinkay J. Effect of mobilization of the anterior hip capsule on gluteus maximus strength. *J Manual Manip Ther.* 2002;10(4):218–224.
53. Makofsky H, Siji P, Abbruzzese J, et al. Immediate effect of grade IV inferior hip joint mobilization on hip abductor torque: a pilot study. *J Manual Manip Ther.* 2007;15(2):103–111.
54. Paquet N, Malounin F, Richards CL. Hip-spine movement interaction and muscle activation patterns during sagittal trunk movements in low back pain patients. *Spine.* 1993;19:596–603.

Copyright of Journal of Sport Rehabilitation is the property of Human Kinetics Publishers, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.